

Information sheet for customer

Tsumura-Kampo Jumihaidokuto [十味敗毒湯] Extract Granules

【Category 2 OTC medicines】

INDICATIONS

Following symptoms in persons with moderate physical strength who have skin disease with redness which is sometimes suppurating:

Early stage of suppurative or acute skin disease, rash, eczema/dermatitis, and athlete's foot.



! CAUTION FOR USE

●Who should seek consultation

1. The following persons should consult a physician, pharmacist, or registered distributor before taking this medicine.
 - (1) Persons undergoing medical treatment from a physician.
 - (2) Pregnant women or women suspected of being pregnant.
 - (3) Persons who is physically weak (declining of strength or constitutional weakness).
 - (4) Persons who have gastrointestinal weakness.
 - (5) Persons who have experienced rash, redness, itching, etc. due to medicines or something.
2. Stop taking this medicine and consult a physician, pharmacist or registered distributor if symptoms do not improve after taking this medicine for about 1 month (about 1 week if you take this medicine for the early phase of suppurative or acute skin disease). Take this leaflet with you.
3. Stop taking this medicine and consult a physician, pharmacist or registered distributor if symptoms may make progress after taking this medicine. Take this leaflet with you.

DOSAGE

Take the following dosages with water or warm water before a meal.

Age	One dosage	Daily dosage
15 years and over	1 packet(1.875 g)	2 times a day
7-14 years	2/3 packet	
4-6 years	1/2 packet	
2-3 years	1/3 packet	
Less than 2 years	Do not use	

[Cautions] This medication should be given to children only under adult supervision

INGREDIENTS

2 packets (3.75 g) of this medicine contain 1.75 g of Jumihaidokuto extract of the following mixed crude drugs;

JP Platycodon Root	1.5 g
JP Bupleurum Root	1.5 g
JP Cnidium Rhizome	1.5 g
JP Poria Sclerotium	1.5 g
JP Quercus Bark	1.5 g
JP Aralia Rhizome	0.75 g
JP Saposhnikovia Root	0.75 g
JP Glycyrrhiza	0.5 g
JP Schizonepeta Spike	0.5 g
JP Ginger	0.5 g

(JP: The Japanese Pharmacopoeia)

Inactive ingredients: JP Magnesium Stearate, JP Lactose Hydrate.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Store in a cool, dry place free from direct sunlight.
- (2) Keep all medication out of the reach of children.
- (3) When you take this medicine in divided dose, fold over the opened edge of the packet and store it.
The remaining medicine should be taken within 2 days.
- (4) Since the medicine contains crude drugs, the granules color may vary for individual products, however this has no influence on the efficacy of this medicine.
- (5) Do not take this medicine after the expiration date.

Production of this product has been approved for domestic sales in accordance with the Pharmaceutical Affairs Law of Japan.



Tsumura&Co. 2-17-11, Akasaka, Minato-ku, Tokyo107-8521, JAPAN.
<http://www.tsumura.co.jp/>