

Information sheet for customer

Tsumura-Kampo Daikenchuto[大建中湯]

Extract Granules

【Category 2 OTC medicines】

INDICATIONS

Following symptoms in persons with weak physical strength who complain of abdominal chills and pain:

Lower abdominal pain or sense of abdominal distension.



! CAUTION FOR USE

●Who should seek consultation

1. The following persons should consult a physician, pharmacist, or registered distributor before taking this medicine.
 - (1) Persons undergoing medical treatment from a physician.
 - (2) Pregnant women or women suspected of being pregnant.
 - (3) Persons who have experienced rash, redness, itching, etc. due to medicines or something.
 - (4) Persons diagnosed as having the following: liver disease.

2. The following symptoms may be adverse reactions to this medication. If any of these symptoms occur, stop taking this medicine immediately and consult a physician, pharmacist, or registered distributor. Take the instruction leaflet with you.

| Related areas | Symptoms |
|---------------|------------------------|
| Skin | Rash, redness, itching |

In very rare cases the following serious symptoms may occur. In this case, consult a physician immediately.

| Symptoms' name | Symptoms |
|--------------------------|---|
| Interstitial pneumonia | When go upstairs or exert yourself too much, symptoms such as shortness of breath, breathing difficulties, dry cough, and fever will appear. They will appear suddenly and persist. |
| Hepatic function failure | Following symptoms appear: fever, itching, rash, jaundice (yellowing of skin and white of eyes), brown urine, whole-body tiredness, loss of appetite. |

3. Stop taking this medicine and consult a physician, pharmacist or registered distributor if symptoms do not improve after taking this medicine for about 1 week. Take this leaflet with you.

DOSAGE

Take the following dosages with water or warm water before a meal.

| Age | One dosage | Daily dosage |
|-------------------|-----------------|---------------|
| 15 years and over | 1 packet(2.5 g) | 3 times a day |
| 7-14 years | 2/3 packet | |
| 4-6 years | 1/2 packet | |
| 2-3 years | 1/3 packet | |
| Less than 2 years | Do not use | |

[Cautions] This medication should be given to children only under adult supervision

INGREDIENTS

3 packets (7.50 g) of this medicine contain 0.625 g of Mukoi-Daikenchuto extract of the following mixed crude drugs and 5.0 g of JP Koi;

| | |
|----------------------|-------|
| JP Processed Ginger | 2.5 g |
| JP Ginseng | 1.5 g |
| JP Zanthoxylum Fruit | 1.0 g |

(JP : The Japanese Pharmacopoeia)

Inactive ingredients: JP Magnesium Stearate, JP Lactose Hydrate.

PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Store in a cool, dry place free from direct sunlight.
- (2) Keep all medication out of the reach of children.
- (3) When you take this medicine in divided dose, fold over the opened edge of the packet and store it.
The remaining medicine should be taken within 2 days.
- (4) Since the medicine contains crude drugs, the granules color may vary for individual products, however this has no influence on the efficacy of this medicine.
- (5) Do not take this medicine after the expiration date.

Production of this product has been approved for domestic sales in accordance with the Pharmaceutical Affairs Law of Japan.



Tsumura&Co. 2-17-11, Akasaka, Minato-ku, Tokyo107-8521, JAPAN.
<https://www.tsumura.co.jp/>